

An Experimentally Realizable Weiss Model for Disorder-Free Glassiness

P. Chandra,¹ M.V. Feigelman,² M.E. Gershenson³ and L.B. Ioffe^{2,3}

¹*NEC Research Institute, 4 Independence Way, Princeton NJ 08540*

²*Landau Institute for Theoretical Physics, Moscow, RUSSIA*

³*Department of Physics, Rutgers University, Piscataway, NJ 08855*

We summarize recent work on a frustrated periodic long-range Josephson array in a parameter regime where its dynamical behavior is *identical* to that of the $p = 4$ disordered spherical model. We also discuss the physical requirements imposed by the theory on the experimental realization of this superconducting network.

The identification of the key features underlying the physics of glass formation remains an outstanding problem of statistical mechanics. A glassy system has a “memory” of its past behavior and does not explore all of phase space; it breaks ergodicity without thermodynamic selection of a unique state thereby defying description by the standard Gibbs methods. Though glass formation in the absence of intrinsic disorder is a widespread phenomenon, it remains poorly understood even at the mean-field level. Because vitrification is a dynamical transition that is not necessarily accompanied by a static one, it remains outside the framework of conventional Landau theory. The crucial links between glasses with intrinsic and self-generated disorder remain an area of active discussion. [1] Recently several microscopic non-random models for glassiness have been proposed; most were studied via mapping to disordered systems. [2] What common/distinct features are characteristic of glasses with and without randomness? How similar/different are glasses with short- vs. long-range interactions? What aspects of existing theoretical models are relevant for experiment? In this Proceeding we summarize our current efforts to address these questions in a periodic model that is both analytically accessible [3,4] and experimentally realizable. [5]




Fig. 1 The phase diagram of the array (insert) where T_G indicates the temperature associated with the dynamical instability discussed in the text, T_S is the speculated equilibrium transition temperature and T_m is the “superheating” temperature where the low-temperature metastable states cease to exist.

The physical system under study is a stack of two mutually perpendicular sets of N parallel thin wires with Josephson junctions at each node (Figure 1) that is placed in an external transverse field H . The classical thermodynamic variables of this array are the $2N$ superconducting phases associated with each wire. In the absence of an external field the phase differences would be zero at each junction, but this is not possible for finite H and the phases are thus frustrated. Here we assume that the Josephson couplings are sufficiently small so that the induced fields are negligible in comparison with H ; we shall return to this topic when discussing the experimental realization of this network. We can therefore describe the array by the Hamiltonian

$$\mathcal{H} = - \sum_{m,n}^{2N} s_m^* \mathcal{J}_{mn} s_n \quad (1)$$

where $\mathcal{J}_{mn}$ is the coupling matrix

$$\hat{\mathcal{J}} = \begin{pmatrix} 0 & \hat{J} \\ \hat{J}^\dagger & 0 \end{pmatrix} \quad (2)$$

with $J_{jk} = \frac{J_0}{\sqrt{N}} \exp(2\pi i \alpha j k / N)$ and $1 \leq (j, k) \leq N$ where $j(k)$ is the index of the horizontal (vertical) wires; $s_m = e^{i\phi_m}$ where the ϕ_m are the superconducting phases of the $2N$ wires. Here we have introduced the flux per unit strip, $\alpha = N H l^2 / \phi_0$, where l is the inter-node spacing and ϕ_0 is the flux quantum; the normalization has been chosen so that T_G does not scale with N .

Because every horizontal (vertical) wire is linked to every vertical (horizontal) wire, the number of nearest neighbors in this model is N and it is accessible to a mean-field treatment. For the same reason, the free energy barriers separating its low-temperature solutions scale with N . This situation is in marked contrast to that in conventional 2D arrays where the coordination number and hence the barriers are low. [6,7] A similar long-range network with positional disorder was studied previously. [8] For $\alpha \gg 1/N$ the latter system displays a spin glass transition which was mapped onto the Sherrington-Kirkpatrick model [9] for $\alpha \gg 1$; in this field regime there is no residual “ferromagnetic” phase coherence between wires. Physically this glassy behavior occurs because the phase differences associated with the couplings J_{jk} acquire random values and fill the interval $(0, 2\pi)$ uniformly for $\alpha \gg 1/N$. More specifically, there will be no commensurability if the sum

$$\sum_{k=1}^N J_{jk} J_{kl}^\dagger = \sum_{k=1}^N \exp 2\pi i \left\{ \frac{\alpha(j-l)}{N} \right\} k \quad (3)$$

is a smooth function of $(j-l)$ where $\{j, l\}$ (k) are indices labelling horizontal (vertical) wires; this will occur only if the expression in curly brackets on the r.h.s. of (3) is *not* an integer for all k , a condition always satisfied for the disordered array. For the periodic case, this situation is realized in the “incommensurate window” $1/N \ll \alpha \leq 1$; here the phase-ordering unit cell is larger than the system size so that the “crystalline” phase is inaccessible. [3] There are thus no special field values where the number of low-temperature solutions are finite, in contrast to the situation for $\alpha > 1$.

In the thermodynamic limit of $N \rightarrow \infty$ (with fixed array area), the high-temperature approach to the glass transition in the periodic array has been studied [3,4] using a modified Thouless-Anderson-Palmer (TAP) method. [10] Here we discuss the qualitative picture that emerges from these results (cf. Fig. 1), referring the interested reader elsewhere for a more detailed quantitative treatment. [3,4] As T approaches T_m^+ , where $T_m^+ \sim T_0 \approx \frac{J\sqrt{N}}{2\sqrt{\alpha}}$, there appear a number of metastable states in addition to the paramagnetic free-energy minimum. There is no dynamical transition at this temperature, indicating that the system is not “trapped”. We speculate that this is because these states are energetically unfavorable and/or they are separated by low barriers. At lower temperatures, as $T \rightarrow T_G^+$, the paramagnetic minimum is “subdivided” into an extensive number of degenerate metastable states separated by effectively infinite barriers, and the system is dynamically localized into one of them. Qualitatively, in the interval $T_m > T > T_G$ there appear many local minima in the vicinity of the paramagnetic state separated by *finite* barriers; these barriers increase continuously and become infinite at $T = T_G$. Each of these minima is characterized by a finite “site magnetization” $m_i = \langle s_i \rangle_T$ where “site” refers to a wire. When $T > T_G$ thermal fluctuations average over many states so that $\langle m_i \rangle \equiv 0$. At $T = T_G$ the system is localized in one metastable state and there is an associated jump in the Edwards-Anderson order parameter, ($q = \frac{1}{N} \sum_i \langle m_i \rangle^2$). The low-temperature phase is characterized by a finite q and by the presence of a memory, $\lim_{t' \rightarrow \infty} \Delta(t, t') \neq 0$ where $\Delta(t, t')$ is the anomalous response. We expect that at $T = T_G$, the metastable states are degenerate and thus there can be no thermodynamic selection. However

at lower temperatures interactions will probably break this degeneracy and select a few states from this manifold; then we expect an $(t \rightarrow \infty)$ equilibrium first-order transition (T_S) which should be accompanied by a jump in the local magnetization. In order to observe this transition at T_S the array must be equilibrated on a time-scale (t_W) longer than that (t_A) necessary to overcome the barriers separating its metastable states; t_A scales exponentially with the number of wires in the array. Thus the equilibrium transition at T_S is observable *only* if $t_W \rightarrow \infty$ *before* the thermodynamic limit ($N \rightarrow \infty$) is taken; in the opposite order of limits only the dynamical transition occurs.

The periodic array thus exhibits a first-order thermodynamic transition *preceded* by a dynamical instability; the glass transition at T_G is characterized by a diverging relaxation time and an accompanying jump in the Edwards-Anderson order parameter. Furthermore the dynamical coupled equations for the response and correlation functions of this *non-random* network in the regime ($1/N \ll \alpha < 1$) are *identical* to those obtained for the $p = 4$ (disordered) spherical Potts model. [11] By contrast the behavior of the disordered array is similar to that of the $p = 2$ case with coinciding static and dynamic transitions. The possible connection between non-random glasses and $p \geq 3$ (disordered) spherical models has been previously suggested in the literature, [12–14] and the periodic array results give support to those conjectures. [4] Furthermore these long-range Josephson arrays can be built in the laboratory, allowing for parallel theoretical and experimental studies of the “simplest spin glasses” that have previously been an abstraction.

Since the experimental realization of these arrays is important, we would like to identify a number of simplifying assumptions inherent in the theoretical treatment discussed here; in order to test its predictions and probe beyond its realm (finite-size effects etc.) the fabricated system must satisfy certain physical requirements. In particular the model has been only considered in the classical limit where quantum mechanical fluctuations of the phase variables are negligible. Furthermore the fields induced by the Josephson currents must always be small compared to H so that it can frustrate the phases effectively. These constraints put some strong restrictions on the choice of array parameters, which we elaborate below.

We begin with the first condition, that of maintaining weak quantum fluctuations. The strength of the latter is controlled by two parameters: the dimensionless normal resistance $e^2 R / \hbar$ and the ratio E_C / E_J where E_C and E_J are the Coulomb and the Josephson energy scales respectively associated with each individual phase variables. The state of any array (or junction) is conveniently described by its position on the plane defined by these dimensionless parameters, E_C / E_J and $e^2 R / \hbar$; the resulting plot is called a Schmid diagram. [15] For a conventional short-range array $R \sim R_0$ and $E_C \sim e^2 / C_0$ where R_0 (C_0) is a resistance (capacitance) of an individual junction. We require that the capacitance of the individual wires and the dissipation in the system should be sufficiently large to suppress the quantum fluctuations of phase and charge effects.

According to the Ambegaokar-Baratoff relation, [16] the normal-state resistance of a single junction in the fabricated long-range arrays should be large ($R_0 \gg \hbar / e^2$) so that the associated screening current is small; for the sake of example we shall assume $R_0 \approx 100 \text{ k}\Omega$ and an individual capacitance of $C \approx 1 \cdot 10^{-15} \text{ F}$, parameters associated with preliminary $Al - Al_2O_3 - Al$ tunnel-junction arrays already fabricated. [5] We note that the Josephson effect for a single tunnel-junction of such a large resistance (or for a short-range array of such junctions) would be completely overwhelmed by quantum fluctuations and Coulomb blockade effects. [17–20] Indeed, it is known that conventional nearest-neighbor arrays exhibit insulating behavior if their individual junction resistivance is more than $15 \text{ k}\Omega$. [19] However in the long-range Josephson arrays all the junctions associated with a particular wire are connected in parallel, so that the effective resistance and capacitance determining their position on the Schmid diagram [15] are $R = R_0 / N$ and $C = N C_0$ respectively. Even for a small array with $N = 100$ the effective resistance and capacitance are $R \sim 1 \text{ k}\Omega$ and $C \sim 1 \cdot 10^{-13} \text{ F}$ (corresponding to the charging energy $E_C \sim 10 \text{ mK}$) respectively so that quantum fluctuations and charging effects are negligible for these networks. We note, however, that if the number of wires (and hence the number of neighbors) were decreased significantly the parameters of the individual junctions would have to be altered to avoid complications of charging effects.

We would also like to minimize the effects of everpresent screening currents to ensure that the external field frustrates the superconducting phases effectively. More specifically we demand that the induced flux in the array is less than a flux quantum

$$L i_{eff} < \Phi_0 \quad (4)$$

so that all fields and phase gradients produced by the diamagnetic currents are negligible. In order to determine the maximum induced flux in the array we need only look at its largest loop; our condition then becomes

$$(Nl) i_{eff} < \Phi_0 \quad (5)$$

where l is the distance between nodes in the array. For the frustrated case, the maximum field occurs when the currents add incoherently in one direction and coherently in the other [8] so that

$$i_{eff} = \left(\sqrt{NN}\right) i_c. \quad (6)$$

where i_c is the critical current associated with an individual junction. Using

$$i_c \sim \frac{2e}{\hbar} E_J = \frac{2e}{\hbar} \left(\frac{k_B T_c}{\sqrt{N}} \right) \quad (7)$$

we find that the necessary condition for weak induced fields becomes

$$E_J \lesssim \frac{\Phi_0^2}{lN^{5/2}}. \quad (8)$$

Thus there is a delicate balance to be obtained between the need for many neighbors (and high free-energy barriers) and the requirement (8) that the effective London penetration length be substantial larger than the system size; unfortunately early experimental attempts to realize these arrays were not able to satisfy these restrictions due to resolution limitations associated with the optical lithography methods used. [21,22] Furthermore the predicted glassy behavior is only observable in the field range $\phi_0/(Nl)^2 < H < \phi_0/l^2$; where the minimum and maximum correspond to a single flux quantum in the whole array and in a single plaquette. Even with the use of the e-beam lithography ($l = 0.5 \mu m$) the requirement (8) can only be satisfied for an array of $Al - Al_2O_3 - Al$ junctions ($E_J = 1.10^{-17} \text{ Erg}$) with the number of wires N less than 300. For these parameters with $N = 100$ $\phi_0/(Nl)^2$ is of the order of 0.01 Oe , and a careful screening of the earth's magnetic field is necessary.

Once the arrays are fabricated to specification what measurements should be performed? The field-dependence of the critical temperature of both disordered and periodic arrays is crucial in establishing the effectiveness of the external field in frustrating the phases. It is also a good test of “phase freezing” into a glassy phase, and the functional dependence $T_c(H) \propto H^{-1/2}$ is expected over a wide field and temperature range for both types of arrays. In order to firmly establish the presence of large barriers imperative for glassiness, the history-dependence of the critical current j_c should be studied (i.e. its dependence on the *path* in the $T - H$ plane) in both arrays. We note that below $T_c(\alpha)$ we expect a faster increase in $j_c(T)$ for the periodic network than for its disordered counterpart; [8] specifics for the former case have yet to be determined. We expect that the diverging relaxation time at T_c can be accessed experimentally via the a.c. response to a time-varying magnetic field $H(t)$; the associated ac susceptibility is

$$\chi_\omega = \frac{\partial M_\omega}{\partial H_\omega} = \frac{C(\alpha, N)\omega}{\omega + i/t_R(T)} \quad (9)$$

where t_R is the longest response of the system that diverges at $T = T_G$ and $C(\alpha, N)$ is to be found elsewhere. [4] The $\omega \rightarrow 0$ limit of the a.c. susceptibility jumps to a finite value at $T = T_G$, indicating the development of a finite superconducting stiffness at the transition. Therefore measurement of this a.c. response in a fabricated array would probe its predicted glassiness.

Work is currently in progress to study the physical properties of the periodic model in its non-ergodic regime ($T < T_G$). [23] We note that the glass transition temperature T_G described above corresponds to the system going out of equilibrium as it is cooled infinitesimally slowly from high temperatures. In practice there will always be a finite-rate of cooling and the effective glass transition will occur when the system drops out of equilibrium. If we define the reduced temperature $\Theta \equiv \left(\frac{T-T_G}{T_G}\right)$ then the effective glass transition will occur when the time associated with cooling, $t_c \approx \frac{\Theta}{d\Theta/dt}$ is equal to the relaxation time at that temperature, $t_R \approx t_0\theta^{-\nu}$ where t_0 and ν are determined from the high-temperature analysis elsewhere; [4] the effective glass transition temperature and the associated time are indicated as $\tilde{T}_G$ and $\tilde{t}_G$ in Figure 2. There is evidence for history-dependence; the system's response is very different if an additional field is turned on at a time, t_H , during or after the cooling process and a subsequent measurement is taken at time t_M (cf. Figure 2), reminiscent of the zero-field cooled vs. field-cooled susceptibility observed in spin glasses. Furthermore an “ageing” time-scale t^* seems to exist in this system (cf. Fig. 2); if the field is turned on during the cooling process at time t_H and a subsequent measurement is taken at time t_M such that $t_H - t_M < t^*$ the system “remembers” the cooling process; otherwise ($t_H - t_M > t^*$) it “forgets” it completely. Preliminary results suggest that t^* has a decreasing temperature-dependence and probably vanishes at low temperatures. This presence of an additional “ageing” time-scale does not exist in the Sherrington-Kirkpatrick theory, though it is indeed observed in experimental spin glasses; it is thus amusing that it emerges from the study of a non-random model.




Fig. 2 Schematics of (a) Temperature (T) (b) Application of an Additional Field (ΔH) vs time (t) indicating the time-scales involved in a finite-cooling experiment as discussed in the text; $\tilde{T}_G$ and $\tilde{t}_G$ refer to the “effective glass transition” when the system goes out of equilibrium, and t_H , t_M and t^* are the time-scales associated with the onset of an additional field, a subsequent measurement and ageing.

In conclusion we have presented a summary of recent work on a periodic model that displays “freezing” into one of an extensive number of metastable states without thermodynamic selection. This glass transition is characterized by a diverging relaxation time and an accompanying jump in the Edwards-Anderson order parameter. At low field strengths the dynamical behavior of this system is identical to that of the $p = 4$ (disordered) spherical model. A key advantage of this system is that it can be built in the laboratory, thereby allowing for the possibility of detailed interplay between theory and experiment. The behavior of a similar array with a reduced number of neighbors is a completely open question that could be important in understanding the relevance of long-range models to real experimental glasses. Finally, since any uncertainty in the position of the wires introduces randomness in the array, the continued study of this network could also offer an opportunity to study the crossover between glasses with spontaneously generated and quenched disorder.

[1] e.g. T.R. Kirkpatrick and D. Thirumalai, PRB **36**, 5388 (1987); J.P. Bouchaud and M. Mezard, J. Phys. I **4**, 1109 (1994); S. Franz and J. Herz, PRL **74**, 2115 (1995); M. Potters and G. Parisi, unpublished (cond-mat preprint 9503009) and

references therein.

- [2] e.g. E. Marinari, G. Parisi and F. Ritort, *J. Phys. A*, **27**, 7647 (1994); M. Mezard and G. Parisi, unpublished.
- [3] P. Chandra, L. B. Ioffe and D. Sherrington, *Phys. Rev. Lett.* **75**, 713 (1995).
- [4] P.Chandra, M.V. Feigelman and L.B. Ioffe, *Phys. Rev. Lett.* **76**, 4805 (1996).
- [5] M.E. Gershenson et al., To Be Published.
- [6] e.g. J.E. Mooij and G.B.J. Schön ed., “Coherence in Superconducting Networks,” *Physica* **152B**, 1 (1988).
- [7] J. P. Carini, *Phys. Rev B* **38**, 63 (1988).
- [8] V. M. Vinokur, L. B. Ioffe, A. I. Larkin, M. V. Feigelman, *Sov. Phys. JETP* **66**, 198 (1987).
- [9] D. Sherrington and S. Kirkpatrick, *Phys. Rev. B* **35**, 1792 (1975).
- [10] D.J. Thouless, P.W. Anderson and R.G. Palmer, *Phil. Mag.*, **35**, 593 (1977).
- [11] A. Crisanti, H. Horner, H.-J. Sommers, *Z. Phys. B*, **92**, 257 (1993).
- [12] T.R. Kirkpatrick and D. Thirumalai, *Phys. Rev. B*, **36**, 5388 (1987); T.R. Kirkpatrick, D. Thirumalai and P.G. Wolynes, *Phys. Rev. B*, **40**, 1045 (1989).
- [13] G. Parisi in J.J. Brey, J. Marro, J.M. Rubi and M. San Miguel, eds. *Twenty Five Years of Non-Equilibrium Statistical Mechanics; Proc. of the Thirteenth Sitges Conference*, (Springer-Verlag, Berlin 1995), pp. 135-42.
- [14] S. Franz and J. Herz, *Phys. Rev. Lett.*, **74**, 2115 (1995).
- [15] A. Schmid, *Phys. Rev. Lett.* **51**, 1506 (1983); R. Fazio and G. Shon, *Phys. Rev. B* **43**, 5307 (1991).
- [16] V. Ambegaoker and A.Baratoff, *Phys. Rev. Lett.* **10**, 486 (1963).
- [17] P. Delsing, C. D. Chen, D. B. Haviland, Y. Harada and T. Cleson, *Phys. Rev. B* **50**, 3959 (1994); J. E. Mooij and G. Shon in “Single Charge Tunneling: Coulomb Blockade Phenomena in Nanostructures” ed. by H. Grabert and M. Devoret, Plenum NY (1992).
- [18] M.Iansiti, A.T.Johnson, C.J.Lobb, and M.Tinkham. *Phys. Rev. Lett.* **60**, 2414 (1988).
- [19] J. E. Mooij, B. J. van Wess, L. J. Geerligs, M. Peters, R. Fazio and G. Shon, *Phys. Rev. Lett.* **65**, 645 (1990).
- [20] M. Tinkham in in “Single Charge Tunneling: Coulom Blocade Phenomena in Nanostructures” ed. by H. Grabert and M. Devoret, Plenum NY (1992).
- [21] L. L. Sohn, M. T. Tuominen, M. S. Rzchowski, J. U. Free, and M. Tinkham. *Phys.Rev.B* **47**, 975 (1993).
- [22] L. L. Sohn, M. S. Rzchowski, J.U.Free, and M.Tinkham. *Phys. Rev. B* **47**, 967 (1993).
- [23] P. Chandra, M.V. Feigelman, L.B. Ioffe and D.M. Kagan, To be published.